

---

Vol. 8 (4): 799-808 (2018)

## **LARGE-SCALE AEROSPACE PHOTOGRAPHY, SOIL-GEOBOTANIC PROFILING IN THE GEOBOTANICAL MAPPING (*DISCOVER ASPECTS OF THE PHYTOCOENOTIC AND BIOGEOCOENOTIC DIVERSITY*)**

Alexander Sizykh<sup>1\*</sup> Victor Voronin<sup>1</sup>, Alexey Shekhovtsov<sup>2</sup>

<sup>1</sup>*Siberian Institute of Plant Physiology and Biochemistry SB RAS, Irkutsk, 664033 Irkutsk, Lermontova str., 132, Russia;*

<sup>2</sup>*V.B. Soshava Institute of Geography SB RAS, Irkutsk, 664033, Ulan-Batorskaya str., 1, Russia;*

\*Corresponding author Alexander Sizykh: [alexander.sizykh@gmail.com](mailto:alexander.sizykh@gmail.com);

Received August, 2018; Accepted September, 2018; Published October, 2018;

DOI: <https://doi.org/10.31407/ijeess8422>

UOI license: <http://u-o-i.org/1.01/ijeess/41633559>

### **ABSTRACT**

The aim of the studies is to reveal phytocoenotic diversity and peculiarities of spatial-temporal organization of phytocoenoses with vegetation types contact and of extrazonal coenoses within zonal taiga in different Pre-Baikalian regions. The objects of studies are phytocoenoses of zonal vegetation types contact as well as of coenoses formed under extrazonal conditions. The subject of studies is to reveal phytocoenotic (and structural-dynamic) peculiarities of environments contact phytocoenoses and to determine the vector of their development on the background of modern climate dynamics.

**Key words:** phytocoenotic diversity, climate dynamics, vegetation types